

JOSLIN DIABETES CENTER and JOSLIN CLINIC
GUIDELINE for INPATIENT MANAGEMENT OF SURGICAL and ICU PATIENTS
(Pre, Peri and Postoperative Care) 4/30/07

The Joslin Clinical Guideline for Inpatient Management of Surgical Patients with Diabetes is designed to assist primary care physicians and specialists to individualize the care and set goals for adult, non-pregnant patients with diabetes who are undergoing surgery. This Guideline focuses on the unique needs of the patient with diabetes. It is not intended to replace sound medical judgment or clinical decision-making and may need to be adapted for certain patient care situations where more or less stringent interventions are necessary.

The objectives of the Joslin Clinical Diabetes Guidelines are to support clinical practice and to influence clinical behaviors in order to improve clinical outcomes and assure that patient expectations are reasonable and informed. Guidelines are developed and approved through the Clinical Oversight Committee that reports to the Joslin Clinic Medical Director of Joslin Diabetes Center. The Clinical Guidelines are established after careful review of current evidence, medical literature and sound clinical practice. This Guideline will be reviewed periodically and modified as clinical practice evolves and medical evidence suggests.

SURGERY ALGORITHM: FOR PATIENTS WITH EXISTING DIABETES

The *Joslin Clinical Guideline for Inpatient Management of Surgical Patients with Diabetes and ICU Patients* uses one formula for “splitting” the insulin; other reasonable formulae exist and are also acceptable.

Aim for Early AM Booking

Day and Evening Prior to Surgery:

- Usual diet and insulin dose (NPH, glargine, detemir, regular, aspart, glulisine, lispro, inhaled insulin, insulin via pump, 70/30, 75/25, or 50/50 insulin) or oral antihyperglycemic medications
- Check blood glucose (BG) at bedtime; if BG > 180 mg/dl, instruct patient to take insulin according to subcutaneous algorithm or per individualized instructions; if hypoglycemic at bedtime or overnight, instruct patient to treat with glucose gel

Morning of Surgery

- If fasting after midnight, give ½ usual dose intermediate (NPH) or full dose long-acting (glargine or detemir) insulin; no change in basal rate for insulin pump patients; no rapid or short-acting insulin; no oral antihyperglycemic medication; no exenatide or pramlintide
- If the patient is coming in from home on pre-mixed insulin (70/30, 75/25, 50/50) and is NPO, less than ½ of the usual morning dose is recommended to avoid hypoglycemia. The optimal regimen would be to give ½ the usual morning dose as NPH insulin.
- If not fasting, give usual dose of insulin
- Check BG every 2 hours before and during surgery; insulin pump patients can maintain basal rate during surgery or be changed to IV insulin infusion or subcutaneous injections to maintain blood glucose target.

Maintenance of Hydration


- During surgery the patient should receive maintenance IV fluids without dextrose (e.g. LR rather than D5LR).
- If an insulin infusion is required, D5W at 40 ml/hr or D10W at 20 ml/hr should be started to provide adequate substrate. This is not required if adequately managed with subcutaneous insulin.
- Patients receiving insulin infusion should receive at least 50 g glucose/24 hours.

Major Surgery	Non-Major Surgery			
E.g., chest or abdominal cavity, LE bypass, transplant, spinal or brain surgery requiring general anesthesia, total hip or knee replacement, surgery anticipated to be > 4 hours <div style="text-align: center;">↓</div> Start IV Insulin (See <i>Insulin Infusion Algorithm</i> pg. 2)	BG < 80 mg/dl ↓ Give at least 100 ml D ₁₀ W IV or 25 – 50 ml (1/2 – 1 amp) of D ₅₀ ↓ Check BG in 15-30 min.	BG 80-100 mg/dl ↓ Begin D ₅ W at 40 ml/hour or D10 W at 20 ml/hour Check BG in 1 hour	BG 101-150 mg/dl ↓ Continue to monitor BG every 2 hours	BG > 150 mg/dl ↓ Begin IV insulin (See <i>Insulin Infusion Algorithm</i> pg. 2) or subcutaneous insulin algorithm

Postoperative Management

- Check BG when patient returns to postanesthesia unit; base frequency on BG during surgery
- Administer insulin according to subcutaneous algorithm or insulin infusion algorithm

- Use maintenance IV fluids without dextrose (e.g. ½ NS rather than D5W ½ NS). If on subcutaneous insulin no additional IV dextrose is required if the patient is not malnourished or in a severely catabolic state. If on an insulin drip substrate must be provided as constant dextrose infusion (e.g. D5W @ 10-40 ml/hr).



PRE, INTRA and POST-OPERATIVE SUBCUTANEOUS SHORT-ACTING INSULIN ALGORITHM

This algorithm can be used a) to supplement an insulin regimen already in place, b) for patients previously on oral antihyperglycemic medications or c) for patients with hyperglycemia without a diagnosis of diabetes. For patients without a diagnosis of diabetes who are normoglycemic prior to surgery, there is no evidence to support a specific frequency of monitoring glucose during surgery. Certain major surgical procedures such as cardiovascular and transplant surgery are associated with hyperglycemia and warrant frequent blood glucose monitoring during and after surgery.

- Monitor glucose level and administer insulin:
 - For glucose level >180 mg/dl, check hourly; if no improvement in glycemic control, consider insulin dosing according to next higher weight class
 - Every 4-6 hours if using regular insulin (short-acting)
 - Every 2-4 hours if using aspart, glulisine, lispro or inhaled insulin (rapid-acting)

	Weight Class I (<175 lbs/80 kg)	Weight Class II (175-220 lbs/81-99 kg)	Weight Class III (>220 lbs/100 kg)
BG (mg/dl)	Insulin Units (subcut)	Insulin Units (subcut)	Insulin Units (subcut)
<150	0 unit	1 unit	2 units
150-180	1 unit	2 units	4 units
181-200	2 units	4 units	6 units
>200	Begin insulin infusion	Begin insulin infusion	Begin insulin infusion

PRE, INTRA and POST OPERATIVE IV INSULIN INFUSION ALGORITHM

Decision to initiate IV insulin:

- If BG >150 mg/dl twice intra-operatively
- If BG >110 mg/dl twice postoperatively for cardiothoracic surgery
- If BG > 150 mg/dl twice in intensive care units in non-cardiothoracic cases

A number of well-validated insulin infusion protocols have been shown to work effectively. There is little data to show any one is superior. Two sample algorithms are provided on pages 3 and 4: one designed to target BG 101-150 mg/dl and another to target BG 80-110 mg/dl.

- If BG < 180 mg/dl, begin D5W at 40 ml/hr or D10W at 20 ml/hr. Maintenance IV fluids without dextrose (e.g. LR or ½ NS or NS) will be added to this in accordance with the patient’s volume requirements. For prevention of ketosis, in most individuals, 50g/24 hours of glucose is generally recommended

Calculating the Initial Insulin Dose
If BG > 180 mg/dl, give stat dose of IV insulin, 0.1 units/kg body weight
For patients having major surgery, larger starting doses can be given: initiate an hourly rate of total daily dose of insulin divided by 24
For patients who have never been on insulin, 0.02 units/kg body weight/hr
For acute surgical patients, e.g., cardiothoracic or transplant, higher starting doses may be necessary
For patients on total parenteral nutrition (TPN), insulin infusion is in addition to insulin currently administered in the TPN solution

Alternative Initial Dose		
Blood glucose (mg/dl)	Regular Insulin (bolus)	Regular Insulin (infusion per hour)
151-200	No Bolus	2 units IV
201-250	3 units IV	2 units IV
251-300	6 units IV	3 units IV
301-350	9 units IV	3 units IV
>350	10 units IV	4 units IV

↓
Check BG Hourly

RECOMMENDED

**INSULIN INFUSION ALGORITHM FOR INTRAOPERATIVE and MEDICAL ICU
(Target BG 101 – 150 mg/dl)**

Insulin dose adjustments using this algorithm do not replace sound medical judgment.

*Whichever is greater change Previous Blood Glucose (mg/dl)

Current Blood Glucose (mg/dl)	Previous Blood Glucose (mg/dl)									
	<60	60-80	81-100	101-150	151-200	201-250	251-300	301-400	>400	
<60	Hold drip and give 1 amp 50% glucose and check BG every 30 minutes until >100 mg/dl and then re-initiate drip at 50% previous rate									
60-80	Hold drip and check BG every 30 minutes until >100 mg/dL and then re-initiate drip at 50% previous rate									
81-100	↓ rate by 1 unit/hr	No change	↓ rate by 25% or 0.5 units/hr*	↓ rate by 50% or 2 units/hr*			↓ rate by 75% or 2 units/hr*			
101-150	No Change					↓ rate by 50% or 2 units/hr*				
151-200	↑ rate by 1 unit/hr		↑ rate by 0.5 units/hr		↑ rate by 25% or 1 unit/hr*	No Change		↓ rate by 25% or 2 units/hr*		
201-250	↑ rate by 25% or 2 units/hr*				↑ rate by 25% or 1 unit/hr*			↑ rate by 1 unit/hr	No Change	
251-300	↑ rate by 33% or 2.5 units/hr*			↑ rate by 25% or 1.5 units/hr*	↑ rate by 25% or 1 unit/hr*	↑ rate by 1 unit/hr	↑ rate by 1.5 units/hr	↑ rate by 25% or 2 units/hr*	No Change	
301-400	↑ rate by 40% or 3 units/hr*									
>400	↑ rate by 50% or 4 units/hr*									

↓
This algorithm assumes hourly BG checks during insulin dose titration.

↓
If BG in desirable range (101-150 mg/dl) for 4 hours can decrease frequency of BG checks to every 2 hours while BG stays in target.
If experiencing unexplained hypoglycemia or hyperglycemia, investigate and correct causative factors.
If there is any significant change in glycemic source (i.e., parenteral, enteral or oral intake), expect to make insulin adjustment.

Common reasons to discontinue insulin infusion:

- Patient tolerating at least 50% of normal oral intake or enteral feedings
- Clinically appropriate to transfer patient to a unit that does not do insulin infusions
- Patient on stable regimen of TPN with most of insulin already in TPN solution

Two hours before discontinuing insulin infusion, initiate alternative glycemic management:

- **For type 1 DM and type 2 DM previously controlled on insulin:** If NPO, initiate basal subcutaneous insulin (glargine, detemir or NPH) at 80% of the insulin administered over the previous 24 hours by insulin infusion. If the patient is taking more than 50% of usual oral or enteral intake, give 50% of insulin dose as basal based on previous 24 hours of insulin infused or 0.25 units/kg and initiate pre-meal bolus and correction dose to maintain BG in target. Another alternative is to resume pre-hospital insulin regimen. Insulin pump patients can resume pump use based on hospital policy.
- **For type 2 DM previously on oral antihyperglycemic agents:** If patient had good diabetes control previous to hospitalization, a return to oral agent therapy may be considered based on postoperative clinical status; if pre-hospital control was poor, plan for discharge on subcutaneous insulin.

Glossary		
AACE – American Association of Clinical Endocrinologists	IV – Intravenous	NS – Normal saline
BG – Blood glucose	LE – Lower extremity	Subcut - subcutaneously
DM – Diabetes mellitus	LR – Lactated Ringers	TPN – Total parenteral nutrition
ICU – Intensive care unit	NPO – Nothing by mouth	

Approved by Clinical Oversight Committee on 4/30/07.

Guideline Task Force: Co-chairs James Rosenzweig, MD and Elaine Sullivan, MS, RN, CDE; Martin J. Abrahamson, MD, Mark Aronson, MD, George Blackburn, MD, PhD, Vasti Broadstone, MD, Amy Campbell, MS, RD, CDE, Roberta Capelson, MS, ANP, Justine Carr, MD, David Feinbloom, MD, Patricia Folcarelli, MS, RN, Michael Howell, MD, Lyle Mitzner, MD, Steven Quevedo, MD, Patricia Samour, MMSc, RD, Marjorie Serrano, RN, Kenneth Snow, MD, Balachundhar Subramaniam, MD

References:

ACE/ADA Task Force on Inpatient Diabetes. American College of Endocrinology and American Diabetes Association consensus statement on inpatient diabetes and glycemic control. *Endocr Pract* 12:4-13, 2006.

Bolk J, van der Ploeg T, Cornel JH, Arnold AE, Sepers J, Umans VA. Impaired glucose metabolism predicts mortality after a myocardial infarction. *Int J Cardiol* 79:207-214, 2001.

Hwaite SS, Godara H, Song HJ, Rock P. No patient left behind: Evaluation and design of intravenous insulin infusion algorithms. *Endocr Pract* 12:72-78, 2006.

Browning LA, Dumo P. Sliding-scale insulin: An antiquated approach to glycemic control in hospitalized patients. *Am J Health Syst Pharm* 61:1611-1614, 2004.

Capes SE, Hunt D, Malmberg K, Gerstein HC. Stress hyperglycemia and increased risk of death after myocardial infarction in patients with and without diabetes: a systematic overview. *Lancet* 355:773-778, 2000.

Carr JM, Sellke FW, Fey M, Doyle MJ, Krempin JA, de la Torre R, Liddicoat JR. Implementing tight glucose control after coronary artery bypass surgery. *Ann Thorac Surg* 80:902-909, 2005.

Clement S, Braithwaite SS, Magee MF, Ahmann A, Smith EP, Schafer RG, Hirsh IB. Management of diabetes and hyperglycemia in hospitals. *Diabetes Care* 27:553-591, 2004.

Egi M, et al. Intensive insulin therapy in postoperative intensive care unit patients. *Am J Respir Crit Care Med* 173: 407-413, 2006.

Furnary AP, Wu Y, Bookin SO. Effect of hyperglycemia and continuous intravenous insulin infusion on outcomes of cardiac surgical procedures: The Portland Diabetic Project. *Endocr Pract* 10 (suppl 2): 21-33, 2004.

Gandhi GY, Nuttall GA, Abel MD, Mullany CJ, Schaff HV, Williams BA, Schrader LM, Rizza RA, McMahon MM. Intraoperative hyperglycemia and perioperative outcomes in cardiac surgery patients. *Mayo Clin Proc* 80:862-866, 2005.

Goldberg PA, Siegel MD, Sherwin RS, Halickman JI, Lee M, Bailey VA, Lee SL, Dziura JD, Inzucchi SE. Implementation of a safe and effective insulin infusion protocol in a medical intensive care unit. *Diabetes Care* 27:461-467, 2004

Haas L. Improving inpatient diabetes care: Nursing issues. *Endocr Pract* 12:56-60, 2006.

Hirsch I. Inpatient diabetes: Review of data from the cardiac care unit. *Endocr Pract* 12:27-34, 2006.

Inzucchi SE. Management of hyperglycemia in the hospital setting. *N Engl J Med* 355:1903-11, 2006.

Malhotra A. Intensive insulin in intensive care. *N Engl J Med* 354:516-518, 2006.

Najarian J, Swavely D, Wilson E, Merkle L, Wasser T, Hesener Quinn A, Urffer S, Young M. Improving outcomes for diabetic patients undergoing vascular surgery. *Diabetes Spectr* 18:23-60, 2005.

Pittas AG, Siegel RD, Lau J. Insulin therapy and in-hospital mortality in critically ill patients: Systematic review and meta-analysis of randomized controlled trials. *J Parenter Enteral Nutr* 30:164-172, 2006.

Schnell O, Schafer O, Kleybrink S, Doering W, Standl E, Otter W. Intensification of therapeutic approaches reduces mortality in diabetic patients with acute myocardial infarction. *Diabetes Care* 27:455-460, 2004.

Swift CS, Boucher JL. Nutrition care for hospitalized individuals with diabetes. *Diabetes Spectr* 18:34-38, 2005.

Swift CS, Boucher JL. Nutrition therapy for the hospitalized patient with diabetes. *Endocr Pract* 12:61-67, 2006.

Van den Berghe G. Insulin vs. strict blood glucose control to achieve a survival benefit after AMI? *Eur Heart J* 26:639-641, 2005.

Van den Berghe G, Wilmer A, Hermans G, Meersseman W, Wouters PJ, Milants I, Van Wijngaerden E, Bobbaers H, Bouillon R. Intensive insulin therapy in the medical ICU. *N Engl J Med* 354: 449-461, 2006.

Van den Berghe G, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med* 345:1359-1367, 2001.

Vanhorebeek I, Langouche L, Van den Berghe G. Intensive insulin therapy in the intensive care unit: Update on clinical impact and mechanisms of action. *Endocr Pract* 12:14-21, 2006.

JOSLIN CLINICAL OVERSIGHT COMMITTEE		
James Rosenzweig, MD - Chairperson	Om Ganda, MD	Kristi Silver, MD
Richard Beaser, MD	David Feinbloom, MD	Susan Sjostrom, JD
Elizabeth Blair, MS, CS-ANP, CDE	John Hare, MD	Kenneth Snow, MD
Patty Bonsignore, MS, RN, CDE	Lori Laffel, MD, MPH	Robert Stanton, MD
Amy Campbell, MS, RD, CDE	Melinda Maryniuk, MEd, RD, CDE	William Sullivan, MD
Cathy Carver, ANP, CDE	Medha Munshi, M	Howard Wolpert, MD
Jerry Cavallerano, OD, PhD	William Petit, MD	Martin Abrahamson, MD (<i>ex officio</i>)